



Assessing Genetic Variability in Eggplant to Determine Their Resource Value and Breeding Potential for Yield and Related Traits

**Laxman L Nandi ^a, Partha Saha ^{b*}, T.K. Behera ^c,
Y.A. Lyngdoh ^d, A.D. Munshi ^a, N.D. Saha ^b,
Ranjith Kumar Ellur ^e, R.S. Pan ^f, Akriti Verma ^e
and B.S. Tomar ^a**

^a Division of Vegetable Science, ICAR-Indian Agricultural Research Institute, New Delhi, India 110012, India.

^b ICAR-National Institute for Research on Commercial Agriculture, Research Station, Dinhata, Cooch Behar, West Bengal 736135, India.

^c ICAR-Indian Institute of Horticultural Research, Bengaluru-560 089, India.

^d ICAR-Central Potato Research Station 5th mile, Peak View Road, Shillong, Meghalaya – 793009, India.

^e Division of Genetics, ICAR-Indian Agricultural Research Institute, New Delhi, India 110012, India.

^f ICAR Research Complex for Eastern Region Research Centre, Ranchi, Jharkhand- 834010, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2025/v28i103033>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/144637>

Original Research Article

Received: 20/07/2025
Published: 26/09/2025

*Corresponding author: E-mail: hortparth@gmail.com;

Cite as: Laxman L Nandi, Partha Saha, T.K. Behera, Y.A. Lyngdoh, A.D. Munshi, N.D. Saha, Ranjith Kumar Ellur, R.S. Pan, Akriti Verma, and B.S. Tomar. 2025. "Assessing Genetic Variability in Eggplant to Determine Their Resource Value and Breeding Potential for Yield and Related Traits". *Journal of Advances in Biology & Biotechnology* 28 (10):23–35. <https://doi.org/10.9734/jabb/2025/v28i103033>.

ABSTRACT

Eggplant is one of the most important vegetables grown in India. The present study was undertaken to study the genetic variability in eggplant for their agro-morphological traits. Sixty accessions of eggplant including released varieties, advanced breeding lines and wild relatives were assessed for 5 morphological descriptors and 6 fruit descriptors. Genetic component viz., GCV, PCV, heritability and genetic advance as per cent over mean were figured out. Genetic diversity study was carried out based on cluster analysis. Principal component analysis (PCA) and Hierarchical cluster analysis (HCA) were carried out for extracting maximum data. A majority of the characters exhibited maximum range of variation as well as many of these characters also displayed high genotypic and phenotypic variance (>20%) except for the fruit length showing modest estimates of PCV (18.73%). The total yield per plant showed positive association with leaf length (0.319), fruit length (0.447), fruit diameter (0.451), average fruits per plant (0.308) and average fruit weight (0.706) at genotypic level at $p=0.01$ significance. The characters which exhibited a positive correlation with yield were found to have a positive direct effect on yield. Based on D^2 values, 60 accessions were assembled into 7 highly distinct clusters. The maximum intra-cluster distance was observed in cluster I (2.537) followed by IV (2.421) and the maximum inter-cluster distance was observed between clusters II and IV (7.062). The first two components of PCA depicted 50.39% of the total variance and AHC categorized the germplasm into four key groups. Thus, it can be presumed by this concept that there is a broad array of variability exists in this crop for almost all the traits which will have ample scope for improvement of eggplant.

Keywords: Eggplant; genetic diversity; correlation; path analysis; growth; yield; PCA; AHA.

1. INTRODUCTION

The genus *Solanum* is the largest and most economically significant among Solanaceae family, comprising approximately 1,400 species that spread all over the world (Ganaie et al., 2018; Motti 2021; Deesh et al., 2025). Eggplant (*Solanum melongena*; $2n=2X=24$) is one of the chief and indigenous solanaceous vegetable crop which has been cultivated in India since ancient time as well as in different continents of the world. The domestication of eggplant from the wild relative *Solanum incanum* L. occurred 4000 years ago in South East Asia based on morphological, and molecular evidence as well as the high fertility of F_1 hybrids with *S. melongena* (Lester & Hasan 1991; Meyer et al., 2012). India is the land of eggplant diversity and has accumulated large genetic diversity in various parts of the country as land races, primitive cultivars and weedy forms especially in peninsular region of Western, Eastern and North Eastern parts of India (Nandi et al., 2021). Eggplant is being extensively grown in an area of approximately 6.83 lakh hectares, with a production and productivity accounting to 129.98 lakh tonnes (Kumar et al., 2022; Jadhav et al., 2025). Eggplant is well-known for its antioxidants property and its fruit extracts proved to have hepatoprotective, anti-carcinoma, anti-microbial, anti-LDL, anti-viral, anti-diabetic, anti-hyperglycemic, anti-hypertensive anti-

hypercholesterolemia and cardio-protective properties (Nwanna et al., 2016; Gürbüz et al., 2018; Yarmohammadi et al., 2021).

The wild species of eggplant provide an enormous reservoir of favourable alleles against many biotic and abiotic stresses and there hasn't been much progress in terms of the overall improvement of eggplant using wild resources (Dempewolf et al., 2017; Nandi et al., 2021). The success of introgression of trait from wild species to cultivated eggplant depends upon their gene pool. In order to identify the important wild species for developing crosses in eggplant improvement, it is necessary to carry out extensive morphological phenotyping of the eggplant germplasm, wild accessions, parental lines, F_1 s, as well as their advanced progenies (Kaushik et al., 2016).

Morphological characterization is the most effective step in the depiction and classification of germplasm which encourages plant breeders in the better use of germplasm (Nand et al., 2018; Nandi et al., 2021; Lyngdoh et al., 2025). Morphological characterization is shown to be effective in assessing the genetic variability and establishing association between different eggplant varietal groups (Sulaiman and Duangjit 2021). Thus, the current research was pursued with intent of characterization of germplasm for various traits to facilitate in the effective selection

of desirable lines for use in breeding program. The outcomes of this research work would help in recognizing superior accessions and support to execute the eventual breeding programme to foster the varietal improvement programs of eggplant in India.

2. MATERIALS AND METHODS

The present research study was undertaken using sixty eggplant accession, including released varieties, advanced breeding lines and wild relatives, which have been collected from different parts of the country and conserved as an active collection at Division of Vegetable Science, ICAR-Indian Agricultural

Research Institute (IARI), New Delhi, (Fig. 1). Seeds were sown in raised nursery bed to raise seedlings. Four weeks old seedlings were transplanted to the main field at a distance of 75 × 60 cm apart. Light irrigation was given immediately after transplanting and thereafter, irrigation was given based on requirement. The plant protection chemicals were sprayed as per standard package of practices. The crop was fertilized with standard doses of fertilizers at 150:100:50 kg NPK/ha. Phenotypic data were recorded for five morphological and six fruit descriptors on five randomly selected plants as per the guidelines on Distinctiveness, Uniformity, and Stability of eggplant (PPV & FRA 2001).



Fig. 1. The morphology of eggplant accessions used in the study. 1. Sel-91-2, 2. 129-5, 3. 190-10-12, 4. Arka Kusamakar, 5. Arka Neelkant, 6. Arka Nidhi, 7. BB-04, 8. BR-112, 9. DB-09, 10. Pusa Kaushal, 11. G-5, 12. G-9, 13. G-10, 14. G-17, 15. G-22, 16. G-27, 17. G-30, 18. G-29, 19. G-44, 20. G-47, 21. G-57, 22. G-60, 23. G-109, 24. G-110, 25. G-111, 26. G-128, 27. G-129, 28. G-130, 29. G-136, 30. G-143, 31. G-164, 32. G-188, 33. G-190, 34. G-203, 35. IC112991, 36. NDB -25, 37. Pant Rituraj, 38. Pant Samrat, 39. Pusa Purple Cluster, 40. Pusa Purple Long, 41. Punjab Sadabahar, 42. Pusa Ankur, 43. Pusa Anupam, 44. Pusa Bhairav, 45. Pusa Bindu, 46. Pusa Kranti, 47. Pusa Purple Round, 48. Pusa Shyamla, 49. Pusa Upkar, 50. Pusa Uttam, 51. *S. aethiopicum*, 52 *S. incanum*, 53. *S. khasianum*, 54. *S. macrocarpon*, 55. *S. sysimbrifolium*, 56. *S. viarum*, 57. *S. integrifolium*, 58. *S. insanum*, 59. DBWSR-195, 60. DBWL-22 (Serial number 1 from left to right upto 60)

For each trait, the means of the estimates were used in all statistical analysis. The data set consisted of a 60×11 matrix, in which the rows displayed the eggplant accessions and columns were average values of the different traits under study. Analysis of variance was performed in accordance with the framework suggested by Panse and Sukhatme (1967). The mean values were distinguished by TUKEY's Honest Significant Difference at a level of $p \leq .0001$. Genetic component viz., Genotypic Coefficient of Variation (GCV), Phenotypic Coefficient of Variation (PCV), heritability, Genetic advance (GA) and genetic advance as per cent over mean were figured out using software SAS ver. 9.3. Correlation coefficient analysis and path coefficient analysis were performed as delineated by Al-Jibouri et al. (1958) and Wright (1921) respectively. Correlation coefficients matrix further visualized with Correlogram generated using package of "corrplot" in R package version 4.0.2. Genetic diversity study was carried out based on cluster analysis as advocated by Mahalanobis (1936) and Rao (1952).

Chemometric data techniques viz., Principal component analysis (PCA) and Hierarchical cluster analysis (HCA) were carried out for extracting maximum data (Cruz et al. 2013). All variables were auto-scaled before deployment of the chemometrics methods. For assessing principal component we used R programme in built function *prcomp* provided in FactoMineR package and factoextra R package is being used to generate a ggplot2-based PCA visualization. For hierarchical cluster analysis, we followed agglomerative clustering method which is the common type of hierarchical clustering used for grouping objects on the basis of their proximity. To perform agglomerative hierarchical cluster analysis in R, we first computed the pairwise distance matrix employing the euclidean distance. The Ward's minimum variance linkage method was used to establish clusters followed by the ggplot2-based dendrogram were produced using factoextra R package for illustration of clusters. The number of cluster formed in AHC is validated with elbow and silhouette methods.

3. RESULTS

3.1 Mean Performance of Eggplant Accessions for Various Morphological Characteristics

The mean performance of the eggplant accessions for 11 traits corresponding to

morphological and yield traits. The characters under observation showed a wide range of variation such as plant height (cm) (27.92-145.48), plant spreading distance (cm) (52.42-169.37), leaf length (cm) (4.08-20.00), leaf width (cm) (3.37-22.87), number of primary branches (3.97-13.66), fruit length (cm) (2.53-21.80), fruit diameter (cm) (1.72-9.03), fruit index (0.82-8.00), average fruits per plant (9.02-60.12), average weight of fruit (g) (12.15-188.00), and yield per plant (kg) (0.56-2.14).

3.2 Genetic Component Analysis

The estimates of genetic component were detailed in (Table 1). High (>20%) GCV and PCV were noted in almost all the characters except for fruit length (cm), which showed moderate PCV estimates (18.73 percent). The highest GCV (52.7%) and PCV (53.73%) were noticed for fruit index while lowest was for fruit length [GCV (20.87); PCV (18.73)].

3.3 Correlation Coefficient

Total yield per plant recorded a positive association (at $p=0.01$) with leaf length (0.319), fruit length (0.447), fruit diameter (0.451), average fruits per plant (0.308), and average weight of fruit (0.706) at the genotypic level (Fig. 2).

3.4 Path Coefficient Analysis

In the present investigation the characters showing a positive correlation with yield also had a positive direct effect on yield itself (Table 2). Among them the average weight of fruit showed to have the highest positive effect on yield at both genotypic (0.936) and phenotypic level (0.981) which was followed by average fruits per plant ($G=0.740$, $P=675$), fruit length ($G=0.324$, $P=0.368$), fruit diameter ($G=0.132$ $P=0.110$) and leaf length ($G=0.101$; $P=0.133$).

3.5 Genetic Divergence Using Multivariate D^2 Analysis

The 60 eggplant accessions were grouped into 7 highly distinct clusters as based on the D^2 values (Table 3). The numbers of accessions in each cluster are given in descending order i.e. Cluster VI (14), V (13), VII (12), III (8), I (5), II (5), and IV (3), respectively. Some of the wild species form a separate cluster such as *S. integrifolium*, *S. xanthocarpum*, *S. incanum*, *S. insanum*, *S. aethiopicum* in cluster I and *S. khasianum*, *S. viarum*, and *S. sisymbriifolium* in cluster IV. The cluster mean values are provided in Table 4

Table 1. Estimates of range, mean, PCV, GCV, heritability, GA and GA as per cent of mean for growth and yield parameters

	Character	Range	Grand Mean	PCV (%)	GCV (%)	Heritability h^2 (%)	GA	GA as per cent of mean
1	Plant height (cm)	27.92-145.48	70.23	31.09	30.63	97.01	36.78	52.79
2	Plant spreading distance (cm)	52.42-169.37	108.7	27.1	26.4	94.93	48.9	45.02
3	Leaf length (cm)	4.08-20.00	13.59	24.66	23.5	90.79	5.34	39.19
4	Leaf width (cm)	3.37-22.87	9.47	30.2	29.38	94.69	4.68	50.04
5	Number of primary branches	3.97-13.66	8.07	28.97	26.21	81.85	3.34	41.49
6	Fruit length (cm)	2.53-21.80	13.17	18.73	20.87	80.53	2.58	34.63
7	Fruit diameter (cm)	1.72-9.03	4.68	26.65	27.48	78.56	1.44	28.24
8	Fruit index	0.82-8.00	3.18	53.73	52.7	94.06	0.79	53.25
9	Average fruits per plant	9.02-60.12	21.8	49.32	47.32	92.06	16.98	79.45
10	Average weight of fruit (g)	12.15-188.00	79.9	49.93	49.67	98.99	68.51	86.49
11	Yield per plant (kg)	0.56-2.14	1.41	30.31	25.94	73.19	0.55	38.91

Table 2. Genotypic and phenotypic path coefficient analysis among growth and yield parameters

Parameters		Plant height	Plant spreading distance	Leaf length	Leaf width	Number of primary branches	Fruit length	Fruit diameter	Fruit index	Average fruit per plant	Average weight of fruit	Yield per plant
Plant height	G	-0.288	-0.007	0.034	-0.008	0.005	-0.013	0.014	0.023	0.018	0.253	0.043
	P	-0.323	-0.007	0.047	-0.008	0.009	-0.012	0.013	0.032	0.022	0.270	0.031
Plant spreading distance	G	-0.083	-0.023	0.010	-0.007	0.007	-0.024	0.018	0.025	-0.002	0.132	0.057
	P	-0.097	-0.022	0.013	-0.007	0.013	-0.029	0.016	0.035	-0.006	0.140	0.053
Leaf length	G	-0.096	-0.002	0.101	-0.055	-0.015	0.117	0.044	-0.009	-0.175	0.363	0.319**
	P	-0.114	-0.002	0.133	-0.060	-0.028	0.143	0.040	-0.014	-0.178	0.399	0.272**
Leaf width	G	-0.025	-0.002	0.062	-0.090	-0.019	-0.015	0.041	0.024	-0.002	0.094	0.086

Parameters		Plant height	Plant spreading distance	Leaf length	Leaf width	Number of primary branches	Fruit length	Fruit diameter	Fruit index	Average fruit per plant	Average weight of fruit	Yield per plant
	P	-0.029	-0.002	0.087	-0.091	-0.037	-0.018	0.037	0.036	-0.002	0.105	0.066
Number of primary branches	G	-0.025	-0.003	-0.026	0.029	0.060	0.022	-0.018	-0.014	0.040	-0.019	0.085
	P	-0.029	-0.003	-0.037	0.034	0.100	0.031	-0.017	-0.019	0.044	-0.020	0.046
Fruit length	G	0.011	0.002	0.037	0.004	0.004	0.324	0.011	-0.098	-0.322	0.418	0.447**
	P	0.011	0.002	0.052	0.004	0.009	0.368	0.007	-0.137	-0.320	0.452	0.391**
Fruit diameter	G	-0.031	-0.003	0.034	-0.028	-0.008	0.026	0.132	0.068	-0.421	0.629	0.451**
	P	-0.039	-0.003	0.048	-0.031	-0.015	0.023	0.110	0.091	-0.418	0.685	0.398**
Fruit index	G	0.051	0.004	0.007	0.017	0.006	0.245	-0.069	-0.130	0.004	-0.059	0.099
	P	0.059	0.005	0.011	0.019	0.011	0.287	-0.057	-0.176	0.001	-0.061	0.075
Average fruit/plant	G	-0.007	0.000	-0.024	0.000	0.003	-0.141	-0.075	-0.001	0.740	-0.637	0.308**
	P	-0.011	0.000	-0.035	0.000	0.007	-0.174	-0.068	0.000	0.675	-0.701	-0.141
Average weight /fruit	G	-0.078	-0.003	0.039	-0.009	-0.001	0.145	0.089	0.008	-0.504	0.936	0.706**
	P	-0.089	-0.003	0.054	-0.010	-0.002	0.169	0.077	0.011	-0.482	0.981	0.621**

Genotypic Residual effect = 0.291; Phenotypic Residual effect = 0.339, ** Indicates significant at $p=0.01$, * Indicates significant at $p=0.05$, Bold and diagonal values indicate direct effect

Table 3. Clustering pattern of eggplant accessions based on fruit growth and yield parameters using D² analysis

Clusters	No of accessions	Genotype name
I	5	<i>S. integrifolium</i> , <i>S. xanthocarpum</i> , <i>S. incanum</i> , <i>S. insanum</i> , <i>S. aethiopicum</i>
II	5	Pusa Kranti, DBGL-164, DBL-60, DBL-143, DBL-188
III	8	Pusa Ankur, Pusa Bhairav, DBL-109, DBR-130, DBL-203, Sel-91-2, BR-112, BB-04
IV	3	<i>S. khasianum</i> , <i>S. viarum</i> , <i>S. sisymbirifolium</i>
V	13	Pusa Anupam, Pusa Kaushal, Pusa Purple Long, Pusa Shyamla, DBWL-22, Pant Samrat, Arka Kusumakar, Arka Nidhi, DB-129-5, Punjab Sadabahar, NDB-25, Arka Neelkant, DB-09
VI	14	Pusa Bindu, Pusa Purple Cluster, DBWSR-195, DBR-5, DBR-27, DBR-30, DBR-57, DBR-110, DBR-128, DBR-129, DBL-153, Pant Rituraj, IC1112991, <i>S. macrocarpum</i>
VII	12	Pusa Purple Round, Pusa Upkar, Pusa Uttam, DBGR-190, DBR-9, DBR-10, DBL-17, DBR-22, DBR-29, DBR-47, DBL-111, Sel-190-10-12




Fig 2. Heat map showing Pearson's correlation coefficients for growth and parameters (Self-correlations are shown in red colour. Positive correlations are shown in red and negative correlations are shown in blue color. The color intensity is proportional to the correlation coefficients. Correlations that are not significant shown in white colour)

which shows that accessions of cluster IV had the highest mean value for plant height (104.91) and average fruit per plant (53.76), Cluster VII accessions recorded the highest mean value for plant spreading distance (127.32), leaf length (16.56), leaf width (11.07) and fruit diameter (6.54). The number of primary branches (9.69) was maximum in cluster I; and cluster II recorded highest mean values for fruit length (19.91), average weight of fruit (126.53) and yield per plant (1.68). Cluster V recorded maximum fruit index estimate (5.53).

3.6 Principal Component Analysis (PCA)

The first component of the PCA had the highest eigen value of 3.36 and accounted for 30.6% of the total variability of the data set. This was followed with an eigen value of 2.17 showing 19.8% in the second PC and 1.54 and

contributing 14% in the third PC of the variance present in the data. The eigen value in the fourth PC was 1.33 and registered 12.10% of the variance in the data. The remaining seven generated PCs (ie PC5 to PC11) yielded successively smaller eigen value (less than 1) not explaining the significant variability in the data set (23.4% total) (Table 5). Variables that are correlated with PC1 and PC2 seem to be the most significant in illustrating the variability of set of data. The most contributing variables for each dimension explained by correlation plot (Fig. 3). The PC1 was positively correlated with variables such as fruit length, average weight of fruit, fruit diameter, yield per plant, leaf length, leaf width, plant height and plant spreading distance arranged in descending order of their contribution. The PC2 is positively correlated with variables such as fruit per plant, fruit index and number of primary branches.

Table 4. Mean values of cluster for growth and yield parameters

Clusters		Plant height	Plant spreading distance	Leaf length	Leaf width	Number of primary branches	Fruit length	Fruit diameter	Fruit index	Average fruit per plant	Average weight of fruit	Yield per plant
I	Mean	55.85	89.91	6.46	5.26	9.69	7.01	3.87	1.77	22.18	51.09	1.17
	SE±	10.75	36.67	2.1	1.34	3.1	4.16	1.29	0.69	5.68	9.24	0.48
II	Mean	92.2	115.6	16.39	9.57	8.3	19.91	4.4	4.68	13.73	126.53	1.68
	SE±	14.96	17.29	2.45	0.25	1.1	1.42	0.47	0.82	1.9	38.74	0.28
III	Mean	58.74	110.81	12.88	8.28	8.35	14.2	5.84	2.54	16.03	106.46	1.67
	SE±	19.06	27.54	1.41	1.13	1.99	1.22	1.07	0.62	3.62	23.5	0.35
IV	Mean	104.91	125.8	12.41	10.06	8.75	3.06	2.13	1.45	53.76	13.79	0.75
	SE±	35.81	25.37	5.55	1.88	2.29	0.74	0.38	0.19	7.07	1.54	0.18
V	Mean	55.55	88.73	13.23	8.39	8.8	17.18	3.18	5.53	25.78	59.12	1.46
	SE±	12.43	12.39	1.81	1.48	2.6	2.73	0.5	1.23	7.86	14.03	0.29
VI	Mean	65.68	110.33	13.81	10.61	6.55	10.72	4.66	2.44	20.76	57.62	1.14
	SE±	16.42	30.56	1.29	3.85	1.78	2.46	0.9	0.87	6.1	13.42	0.27
VII	Mean	84.54	127.32	16.56	11.07	7.79	13.45	6.54	2.11	15.61	116.43	1.67
	SE±	12.34	30.06	1.88	2.07	1.42	1.32	1.17	0.34	5.4	36.2	0.22

Table 5. Eigen values, % of Variance and Cumulative % of 11 growth and yield parameters

Principal Components	Eigen value	Variance (%)	Cumulative variance (%)
Dim.1	3.36980605	30.6346005	30.63460
Dim.2	2.17384831	19.7622573	50.39686
Dim.3	1.54041458	14.0037689	64.40063
Dim.4	1.33180026	12.1072751	76.50790
Dim.5	0.77105324	7.0095749	83.51748
Dim.6	0.72096336	6.5542123	90.07169
Dim.7	0.60380339	5.4891217	95.56081
Dim.8	0.20361721	1.8510656	97.41188
Dim.9	0.17004225	1.5458386	98.95771
Dim.10	0.08688027	0.7898206	99.74754
Dim.11	0.02777108	0.2524644	100.00000




Fig. 3. Variable correlation plot on the first two principal components




Fig. 4. Agglomerative hierarchical clustering (AHC) of sixty eggplant accessions using Euclidean distance and ward linkage method

3.7 Agglomerative Hierarchical Cluster Analysis (HCA)

Agglomerative Hierarchical Cluster Analysis (HCA) was carried out on the data and grouping

of accessions was done on the basis of similarities. The method of Euclidean distance and Ward Linkage method was used to perform HCA. The results acquired by implying AHC are represented in a dendrogram (Fig. 4). The

dendrogram revealed a four distinct cluster of 60 eggplant accessions based similarities for agro morphological traits. This number of cluster generated in AHC is also validated and confirmed by elbow and silhouette methods. Cluster A constituted of 24 accessions, which include cultivated accessions as well their counterpart such as *S. incanum*, *S. macrocarpum*, *S. integrifolium*, *S. aethiopicum*, *S. xanthocarpum*, and *S. insanum*. Cluster B constituted of 14 accessions. Cluster C constituted 3 accessions, *S. khasianum*, *S. viarum*, and *S. sisymbirifolium*. Cluster D consisted of 19 accessions which all are cultivated types.

4. DISCUSSION

For the accomplishment of any crop breeding programs, plant genetic resources are regarded as the cornerstone. Greater genetic diversity in the germplasm for different traits of interest viz., yield, quality and resistance to biotic and abiotic stress will ensure better chances of progress for these traits. A broad range of variation was ascertained, for many of the traits in the present study which attributed to the existence of adequate quantities of variation among the accessions for the characteristics being studied.

The combination of high heritability, the intensity of selection and amount of variability exist in germplasm influences the genetic gains to be obtained from selection. The traits such as plant height, plant spreading distance, leaf length, leaf width, number of primary branches, fruit length, fruit diameter, fruit index, average fruit weight, number of fruits per plant and total yield per plant recorded significantly high heritability (>60 %) along with high GAM (>20 %). The discrepancies between PCV and GCV estimates were minimal for most of the characters considered which stipulates that the variability present in the trait are presumably due to genetic factors and are comparatively stable. However, the environment has played a minor role in the manifestation of this phenotype. Thus, these traits can be relied upon for further improvement through direct, simple selection based on phenotype. Moderate estimates of PCV were observed for fruit length and for fruit diameter, GCV was higher than PCV. Similar findings have been reported by Devaraju et al. (2020); Lateef et al. (2022). The results obtained from this study suggested that environment played a little role for manifestations of the results. However, a large difference existed among GCV and PCV for characters like

fruit length and yield per plant indicating the role of the environment over this character.

The development in crop yield will greatly depend on the amount of variability present in the breeding materials and the extent to which the major yield contributing characters are inherited from generation to generation which gives us an option for selecting suitable parents for further crop improvement work. High heritability (>60%) estimates along with high GAM (>20%) was estimated for the most of the traits under analysis, which indicates that, additive component is predominant and hence, direct selection would be more effective in improving these traits. Our results are in accordance with reports by Jayalakshmi et al. (2020); Jyoti et al. (2025). Pearson's correlation coefficient is a statistical approach which quantifies direction and magnitude of the association, or "coherence", between two variables. Our results from this research work were aligned with the works reported by Onyia et al. (2020); Kumar et al. (2020) and Lyngdoh et al. (2025). The average fruit per plant has a positive relationship with total yield per plant in genotypic correlation, but it has shown negative association in phenotypic correlation, and this is primarily due to the interaction between the genotype and the environment.

Mahalanobis D^2 statistics is statistical tool based on second order statistics, appropriate for deciding biologically distinct parents in the hybridization programme. Mahalanobis D^2 statistics is successful in grouping 60 accessions into 7 highly distinct clusters based on D^2 values. A detailed analysis suggests that neither of the clusters had all the desirable characteristics, but many important desirable traits are acquired throughout different clusters which can be assembled effectively via crossing the appropriate accessions depending on the breeding priorities. The present study of genetic diversity provides an insight for genetic interrelationships among the accessions of eggplant and demonstrates the presence of sufficient quantities of genetic variation in the present breeding material.

Principal Component Analysis (PCA) is a dimensionality reduction approach which has a profound applications to data reduction for eg., investigative data analysis and genetic diversity of plants. The PCA showed that out of the 11 components, the first four PCs (>1 eigenvalue) explaining about 76.50% of the total variability

present in the 60 eggplant accessions collection was assessed for different growth and yield traits. The rest of the seven components exhibited just 23.5% of the total morphological diversity among the accessions studied. Our findings are endorsed with assertion by Nandi et al. (2018). PCA recognized only a few characters played important role in classifying the variation present in the germplasm. Thus variation in the germplasm accessions cannot be explained based on only few characters hence further Agglomerative Hierarchical Cluster Analysis (HCA) were used to support the Hierarchical Clustering on Principal Components to provide a useful means for estimating morphological diversity within and between the germplasm collections. Similar finding was made by Dhatt et al. (2017).

AHC generated four cluster. Cluster A constituted of 24 accessions, which include cultivated accessions as well their counterpart such as *S. incanum*, *S. macrocarpum*, *S. integrifolium*, *S. aethiopicum*, *S. xanthocarpum*, and *S. insanum*. The rationale for their inclusion along with cultivated accessions is because they are semi domesticated share most similarities with cultivated types for morphological traits and are used for culinary purposes in many other countries. Cluster B constituted of 14 accessions. Cluster C constituted 3 accessions, *S. khasianum*, *S. viarum*, and *S. sisymbriifolium*, which are weedy types and morphologically contrast to cultivated forms. Cluster D consisted of 19 accessions, which are all cultivated types. Genetic divergence is assessed by multivariate analysis in eggplant which helps to select superior accessions in a breeding program to breed high yielding variety.

5. CONCLUSION

The results conclude that a greater amount of genetic diversity exists among germplasm accessions for all the characters studied. For most traits, high GCV and PCV registered, and the disparities among PCV and GCV values were marginal, signaling little implications of environment on trait expression. Selecting such traits would be effective in enhancing the yield potential. Recombination breeding among accessions belonging to cluster having maximum inter cluster distance are likely to achieve heterotic effect and transgressive segregations. Emphasis should be given to the yield components which showed highly significant correlation as well as direct and indirect effects

on yield per plant for further improvement. Accessions from different clusters that have been known for a particular trait may be used as a parent for breeding program with intent of enhancing the specific traits.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENTS

Authors would like to express sincere gratitude to Indian Council of Agricultural Research, New Delhi and Division of Vegetable Science, ICAR-Indian Agricultural Research Institute, New Delhi for providing financial assistance and all the facilities for conducting the experiment under inhouse project (CRSICIARISIL2014017249).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Al Jibouri, H. A., Miller, P. A., & Robinson, H. F. (1958). Genotypic and environmental variances and covariances in an upland cotton cross of interspecific origin. *Agronomy Journal*, 50, 633–636.
- Cruz, A. G., Cadena, R. S., Alvaro, M. B., Sant'Ana, A. S., Oliveira, C. A., & Faria, J. A. et al. (2013). Assessing the use of different chemometric techniques to discriminate low-fat and full-fat yogurts. *LWT - Food Science and Technology*, 50, 210–214.
- Deesh, A. D., Ubaub, L. T., Furlong, M., & Lomavatu, M. F. (2025). Weed population dynamics and composition in a tropical island eggplant (*Solanum melongena*) agroecosystem: Implications for sustainable weed management. *Pacific Science*, 78(4), 429-443.
- Dempewolf, H., Baute, G., Anderson, J., Kilian, B., Smith, C., & Guarino, L. (2017). Past and future use of wild relatives in crop breeding. *Crop Science*, 57, 1070–1082.
- Devaraju, K. M., Srinivasa, V., Lakshmana, D., Singh, T. H., Devappa, V., & Venugopalan, R. (2020). Studies on variability

- components for yield and its attributes in brinjal (*Solanum melongena*) under hill zone of Karnataka. *Journal of Pharmacognosy and Phytochemistry*, 9, 2200-2203.
- Dhatt, A. S., Kaur, G., Sidhu, M. K., & Kaur, S. (2017). Morphological and molecular characterization of *Solanum melongena* and related species. *Vegetable Science*, 44(1), 1-11.
- Ganaie, M. M., Raja, V., Reshi, Z. A., & Verma, V. (2018). Family *Solanaceae*: Taxonomy and modern trends. *Annals of Plant Science*, 7(9), 2403-2414.
- Gürbüz, N., Uluişik, S., Frary, A., Frary, A., & Doğanlar, S. (2018). Health benefits and bioactive compounds of eggplant. *Food Chemistry*, 268, 602-610.
- Jadhav, V. C., Tated, R. G., Bhangale, J. H., Sancheti, S. D., & Gore, K. B. (2025). A review on vegetable seedlings transplantation systems and their importance in Indian agriculture. *International Journal of Vegetable Science*, 1-34.
- Jayalakshmi, K., Praneetha, S., & Poobalan, V. (2020). Genetic variability studies in Tamilnadu land races of brinjal (*Solanum melongena*). *Journal of Pharmacognosy and Phytochemistry*, 9, 2135-2138.
- Jyoti, Singh, D., Dhankhar, S. K., Saini, K. S., Arora, I., Kumar, V., & Pahal, P. (2025). Genetic diversity among brinjal genotypes grown in North-West India. *Journal of Experimental Agriculture International*, 47(5), 700-708.
- Kaushik, P., Prohens, J., Vilanova, S., Gramazio, P., & Plazas, M. (2016). Phenotyping of eggplant wild relatives and interspecific hybrids with conventional and phenomics descriptors provides insight for their potential utilization in breeding. *Frontiers in Plant Science*, 7, 677.
- Kumar, A., Kumar, R., Ansar, M., Akhtar, S., Adarsh, A., & Kumar, V. (2020). Screening against Phomopsis blight in brinjal (*Solanum melongena*). *International Journal of Current Microbiology and Applied Sciences*, 9, 223-228.
- Kumar, T., Peter, K. V., & Sangeetakutty, M. (2022). Seven decades of vegetables research in India. *International Journal of Innovative Horticulture*, 11(2), 135-157.
- Lateef, A., Makhdoomi, M. I., Bashir, T., Hamid, I., & Farooq, S. (2022). Cluster analysis and percent contribution of individual traits towards total genetic divergence in brinjal. *The Pharma Innovation*, 11(11), 231-235.
- Lester, R. N., & Hasan, S. M. Z. (1991). Origin and domestication of the brinjal eggplant, *Solanum melongena*, from *S. incanum*, in Africa and Asia. In J. G. Hawkes, R. N. Lester, M. Nee, & N. Estrada (Eds.), *Solanaceae III: Taxonomy, chemistry, evolution* (pp. 369-387). The Linnean Society of London.
- Lyngdoh, Y. A., Saha, P., Tomar, B. S., Bhardwaj, R., Nandi, L. L., Srivastava, M., Gurung, B., Ranjan, J. K., & Chaukhande, P. (2025). Unveiling the nutraceutical potential of indigenous and exotic eggplant for bioactive compounds and antioxidant activity as well as its suitability to the nutraceutical industry. *Frontier in Plant Science*, 16, 1451462. doi: 10.3389/fpls.2025.1451462.
- Mahalanobis, P. C. (1936). On the generalized distance in statistics. *Proceedings of the National Institute of Sciences (Calcutta)*, 2, 49-55.
- Meyer, R. S., Karol, K. G., Little, D. P., Nee, M. H., & Litt, A. (2012). Phylogeographic relationships among Asian eggplants and new perspectives on eggplant domestication. *Molecular Phylogenetics and Evolution*, 63, 685-701.
- Motti, R. (2021). The *Solanaceae* family: Botanical features and diversity. In *The wild solanums genomes* (pp. 1-9). Springer International Publishing.
- Nand, N., Adarsh, A., Kumar, A., Akhtar, S., Kumar, R., & Ray, P. K. (2018). Morphological characterization of different genotypes of brinjal (*Solanum melongena*). *International Journal of Current Microbiology and Applied Sciences*, 7(01), 2218-2226. <https://doi.org/10.20546/ijcmas.2018.701.267>
- Nandi, L. L., Saha, P., Behera, T. K., Lyngdoh, Y. A., Munshi, A. D., & Saha, N. D. et al. (2021). Genetic characterization and population structure analysis of indigenous and exotic eggplant (*Solanum* spp.) accessions using microsatellite markers. *The Journal of Horticultural Science and Biotechnology*, 96, 73-86.
- Nwanna, E. E., Ibukun, E. O., & Oboh, G. (2016). Effect of some tropical eggplant fruits (*Solanum* spp.) supplemented diet on diabetic neuropathy in experimental male Wistar rats in-vivo. *Functional Foods in Health and Disease*, 6, 661-676.

- Onyia, V. N., Chukwudi, U. P., Ezea, A. C., Atugwu, A. I., & Ene, C. O. (2020). Correlation and path coefficient analyses of yield and yield components of eggplant (*Solanum melongena*) in a coarse-textured Ultisol. *Information Processing in Agriculture*, 7, 173-181.
- Panse, V. G., & Sukhatme, P. V. (1967). *Statistical methods for agricultural workers* (ICAR Publication). New Delhi.
- PPV and FR Act. (2001). Guidelines for the conduct of test for distinctiveness, uniformity and stability on brinjal/eggplant. <http://www.plantauthority.gov.in/pdf/ingazette.pdf>
- Rao, C. R. (1952). *Advanced statistical methods in biometric research*. John Wiley and Sons Inc.
- Sulaiman, N. N. M., & Duangjit, J. (2021). Genetic diversity of eggplant germplasm based on agro-morphological traits under open field and glasshouse conditions (Doctoral dissertation, Kasetsart University).
- Wright, S. (1921). Systems of mating. I. The biometric relations between parent and offspring. *Genetics*, 6, 111.
- Yarmohammadi, F., Rahbardar, M. G., & Hosseinzadeh, H. (2021). Effect of eggplant (*Solanum melongena*) on the metabolic syndrome: A review. *Iranian Journal of Basic Medical Sciences*, 24(4), 420-427. <https://doi.org/10.22038/ijbms.2021.50276.11452>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://pr.sdiarticle5.com/review-history/144637>